

## Tryptic soy agar

REF.	Pack size
1414 001	100 gm
1414 002	500 gm

### Intended Use

Tryptic Soy Agar is used for the cultivation of a wide variety of microorganisms such as Salmonella Typhi isolated from faeces.

### Background

TSA, a general purpose medium, is commonly referred to as Soybean-Casein Digest Agar USP 23. TSA is a nutritious base, and a variety of supplements can be added to enhance this medium. The addition of 5% sterile, defibrinated sheep, horse, or rabbit blood provides an excellent non-selective medium, used to determine hemolytic reactions of bacteria. TSA supplemented with lecithin and Tween 80 is widely used in environmental monitoring. TSA is used in the differentiation of Haemophilus species (the X and V factors are omitted from this medium), and widely used for blood cultures. TSA is commonly used as a maintenance medium for culture collections, and testing bacterial contaminants in cosmetics.

### Principle

Enzymatic Digest of Casein and Enzymatic Digest of Soybean Meal provide the nitrogen, vitamins and carbon in TSA. Sodium Chloride maintains osmotic balance in the medium. Agar is the solidifying agent.

Components	gm/Liter
Sodium chloride	5.0
Enzymatic digest of Casein	15.0
Enzymatic digest of Soybean	5.0
Agar	15.0

Final pH (at 25°C) 7.3 ± 0.2










### Preparation, Storage and Stability

Store the dehydrated medium at 10-30°C and use before the expiry date on the label. Store the prepared medium at 2-8°C. After the desired amount has been taken out, replace the cap tightly to protect from hydration.

### Procedure

- Suspend 40 g of the powder in 1 L distilled water and mix well.
- Heat with frequent agitation and boil for one minute to dissolve the powder completely.
- Sterilize by autoclaving at 121°C for 15 minutes.

### SYMBOLS IN PRODUCT LABELLING

	Authorised Representative		Temperature Limitation
	For in-vitro diagnostic use		Use by/Expiration Date
	Batch Code/Lot number		CAUTION. Consult instructions for use
	Catalogue Number		Manufactured by
	Consult instructions for use		

### Quality Control

#### Appearance

1-Dehydrated Appearance : Beige coloured, homogeneous, free flowing powder.

2- Prepared Appearance : slight hazy and yellow beige in color.

3- Cultural Response : after 18-24 hours at 30-35°C or 35± 2°C for clinical specimens

#### Organisms (ATCC)

<i>Aspergillus niger</i>	Good
<i>Salmonella Typhi</i>	Good
<i>Candida albicans</i>	Good
<i>Pseudomonas aeruginosa</i>	Good
<i>Streptococcus pyogenes</i>	Good

#### Growth

### Interpretation of the results

- Count the number of colonies on the dipstick. Multiply by dilution factor to convert the count to CFU per ml of the sample.
- Contaminant bacteria usually appear in low numbers, which vary in colony morphology

### Precautions

1-Due to nutritional variation, some strains may be encountered that grow poorly or fail to grow on this medium.

### Bibliography

- United States Pharmacopeial Convention. 2007. The United States pharmacopeia, 31st ed., Amended Chapters 61, 62, 111. The United States Pharmacopeial Convention, Rockville, MD.
- Directorate for the Quality of Medicines of the Council of Europe (EDQM). 2007. The European Pharmacopoeia, Amended Chapters 2.6.12, 2.6.13, 5.1.4, Council of Europe, 67075 Strasbourg Cedex, France.
- Japanese Pharmacopoeia. 2007. Society of Japanese Pharmacopoeia. Amended Chapters 35.1, 35.2, 7. The Minister of Health, Labor, and Welfare.
- Baron, E.J., L.R. Peterson, and S.M. Finegold. 1994. Bailey & Scott's diagnostic microbiology, 9th ed. Mosby-Year Book, Inc., St. Louis.

 Spectrum For Diagnostic Industries - Free Zone  
Ismailia Free Zone Industrial Area, Block 5 .  
Cairo- Port said Avenue.  
Ismailia, Egypt  
Tel: +2 064 3488 013 - +2 064 3488 014 Fax: +2 064 3488 015  
www.sdi-fz.com

 MDSS GmbH  
Schiffgraben 41  
30175 Hannover, Germany



IFUFBC05

Rev.(2),18/6/2019

